

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
11 April 2002 (11.04.2002)

PCT

(10) International Publication Number
WO 02/28433 A2(51) International Patent Classification⁷: **A61K 45/08**,
31/421, 31/426, A61P 1/06, 19/02, 21/00, 29/00, 37/00

(21) International Application Number: PCT/GB01/04370

(22) International Filing Date: 2 October 2001 (02.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0024362.6 5 October 2000 (05.10.2000) GB(71) Applicant (for all designated States except US): **GLAXO
GROUP LIMITED** [GB/GB]; Glaxo Wellcome House,
Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BUCHAN, Kevin,**
William [GB/GB]; c/o GlaxoSmithKline, Gunnels Wood
Road, Stevenage, Hertfordshire SG1 2NY (GB).(74) Agent: **GIDDINGS, Peter, John**; GlaxoSmithKline, Cor-
porate Intellectual Property (CN9.25.1), 980 Great West
Road, Brentford, Middlesex TW8 9GS (GB).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,
ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

— without international search report and to be republished
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 02/28433 A2

(54) Title: MEDICAMENTS

(57) Abstract: Methods of prevention or treatment of diseases or conditions where inhibition of NO synthase and/or TNF is desir-
able, the use of PPAR delta activators in such methods and methods for the identification of compounds useful in such treatment.

94.1.1 变更

Medicaments

The present invention is concerned with medicaments. More particularly, the invention relates to compounds that bind to and activate PPAR delta. In another aspect, the present invention relates to methods for prevention or treatment of diseases and conditions mediated by iNOS and/or TNF and methods for identifying compounds useful in the treatment of diseases and conditions mediated by iNOS and/or TNF.

5 Nitric oxide is the endogenous stimulator of the soluble guanylate cyclase enzyme and is involved in a number of biological actions. Excess nitric oxide production is also thought to be involved in a number of conditions, including septic shock and many inflammatory diseases. The biochemical synthesis of nitric oxide from L-arginine is catalysed by the enzyme NO synthase. Many inhibitors of NO synthase have been described and proposed for therapeutic use. More recently, it has been an object in this field to provide NO synthase inhibitors displaying selectivity for either inducible NO synthase (iNOS) or neuronal NO synthase (nNOS) over endothelial NO synthase (eNOS).

10 Peroxisome proliferator activated receptor (hereinafter referred to as PPAR) is a known member of the steroid/retinoid/thyroid hormone receptor family of ligand activated transcription factors and is activated, inter-alia, by high micromolar concentrations of certain peroxisome proliferators. Peroxisome proliferator activated receptor alpha (herein after referred to as PPAR α), peroxisome proliferator activated receptor gamma (hereinafter referred to as PPAR δ) and peroxisome proliferator activated receptor delta (hereinafter referred to as PPAR δ) have respectively been identified as subtypes of PPARs.

25 Tumor necrosis factor (TNF) is known to mediate many biological responses in vivo. Clinical and Pre-clinical studies in animals and humans with specific TNF neutralising antibodies, soluble TNF receptor constructs and TNF detection techniques have implicated TNF as a mediator in numerous pathologies including inflammatory/auto immune diseases or conditions.

The present inventors have surprisingly found that activators of the peroxisome proliferator activated receptor delta (hereinafter referred to as PPAR δ) inhibit iNOS and TNF formation and are therefore useful in the prophylaxis and treatment of conditions for which an inhibitor of NO synthase is indicated, in particular, an inhibitor of iNOS.

The present invention therefore provides the use of PPAR δ activators in the treatment of diseases and conditions for which an inhibitor of NO synthase and/or TNF is indicated (in particular, an inhibitor of iNOS) and methods of treating NOS and/or TNF mediated diseases employing PPAR δ agonists. More particularly, the present invention provides the use of potent and selective PPAR δ agonists for use in the treatment of conditions for which an inhibitor of NO synthase and/or TNF is indicated, (in particular, an inhibitor of iNOS).

15 Description of Drawings

Figure 1: Shows the effect of PPAR delta agonists on lipopolysaccharide (LPS) induced iNOS activity.

20 Figure 2: Shows the effect of a PPAR delta agonist on LPS elevation of iNOS mRNA.

Figure 3: Shows the effect of a PPAR delta agonist on LPS-induced TNF α expression compound to LPS-induced iNOS activation.

25 Conditions for which an inhibitor of NO synthase and/or TNF is indicated, include inflammatory conditions, shock states, immune disorders, and disorders of gastrointestinal motility. PPAR delta agonists and pharmaceutically acceptable salts, solvates, and physiologically functional derivatives thereof may also be of use in the prophylaxis and treatment of diseases of the central nervous system including migraine.

35 By shock states is meant those resulting from overproduction of NO, such as septic shock, haemorrhagic shock, traumatic shock, or shock caused by fulminant hepatic failure or by therapy with cytokines such as TNF, IL-1 and IL-2

or therapy with cytokine-inducing agents, for example 5,6-dimethylxanthenone acetic acid.

5 Examples of inflammatory conditions and immune disorders include those of the joint, particularly arthritis (e.g. rheumatoid arthritis, osteoarthritis, prosthetic joint failure), or the gastrointestinal tract (e.g. ulcerative colitis, Crohn's disease, and other inflammatory bowel and gastrointestinal diseases, gastritis and mucosal inflammation resulting from infection, the enteropathy provoked by non-steroidal anti-inflammatory drugs), of the lung (e.g. adult respiratory distress syndrome, 10 asthma, cystic fibrosis, or chronic obstructive pulmonary disease), of the heart (e.g. myocarditis), of nervous tissue (e.g. multiple sclerosis), of the pancreas, (e.g. inflammation associated with diabetes melitus and complications thereof of the kidney (e.g. glomerulonephritis), of the skin (e.g. dermatitis, psoriasis, eczema, urticaria and burn injury), of the eye (e.g. glaucoma) as well as of 15 transplanted organs (e.g. rejection) and multi-organ diseases (e.g. systemic lupus erythematosus), inflammatory sequelae of viral or bacterial infections and inflammatory conditions associated with atherosclerosis and following hypoxic or ischaemic insults (with or without reperfusion), for example in the brain or in ischaemic heart disease.

20

Disorders of gastrointestinal motility include ileus, for example post-operative ileus and ileus during sepsis.

25

By diseases of the central nervous system is meant those for which overproduction of NO and/or TNF is implicated, for example migraine, psychosis, anxiety, schizophrenia, sleep disorders, cerebral ischaemia, CNS trauma, epilepsy, multiple sclerosis, AIDS dementia, chronic neurodegenerative disease (e.g. Lewy Body Dementia, Huntington's disease, Parkinson's disease, or Alzheimer's disease) and acute and chronic pain, and conditions in which non-adrenergic non-cholinergic nerve may be implicated such as priapism, obesity and hyperphagia.

30

35

Examples of acute pain include musculoskeletal pain, post operative pain and surgical pain. Examples of chronic pain include chronic inflammatory pain (e.g. rheumatoid arthritis and osteoarthritis), neuropathic pain (e.g. post herpetic

上記の疾患を列挙してあるが、各疾患との関係は明確に説明していない。

neuralgia, diabetic neuropathies associated with diabetes, trigeminal neuralgia, pain associated with functional bowel disorders, e.g. irritable bowel syndrome, non cardiac chest pain and sympathetically maintained pain) and pain associated with cancer and fibromyalgia.

Furthermore, inhibition of NO synthase and/or TNF may be of advantage in preventing the lymphocyte loss associated with HIV infection, in increasing the radiosensitivity of tumours during radiotherapy and in reducing tumour growth, tumour progression, angiogenesis, and metastasis.

Preferably, the PPAR delta activators are PPAR δ agonists. As used herein, by "agonist", or "activating compound", or "activator", or the like, is meant those compounds which have a pKi of at least 6.0 to the relevant PPAR, for example human PPAR δ , in the binding assay described below, and which achieve at least 50% activation of the relevant PPAR relative to the appropriate indicated positive control in the transfection assay described, for example in WO 00/08002 at concentrations of 10^{-5} M or less. More preferably, the compounds of this invention achieve 50% activation of human PPAR δ in the transfection assay at concentrations of 10^{-7} M or less.

Most preferably, PPAR delta agonists are selective hPPAR δ agonists. As used herein, a "selective hPPAR δ agonist" is a hPPAR δ agonist whose EC50 for PPAR δ is at least 10 fold lower than its EC50 for PPAR λ and PPAR α . Such selective compounds may be referred to as "10-fold selective". EC50 is defined in the transfection assay described, for example in WO 00/08002 and is the concentration at which a compound achieves 50% of its maximum activity. Most preferred compounds are greater than 100-fold selective hPPAR δ agonists.

The following are further particular aspects of the present invention:

- (a) a PPAR delta agonist (e.g. a compound of formula (I)) for use as a therapeutic agent in the treatment of conditions for which an inhibitor of NO synthase and/or TNF is indicated, in particular, an inhibitor of iNOS;

(b) pharmaceutical formulations comprising a PPARdelta agonist (e.g. a compound of formula (I)) and at least one pharmaceutical carrier, wherein the PPAR delta agonist is present in an amount effective for use in the treatment of conditions for which an inhibitor of NO synthase and/or TNF is indicated, in particular, an inhibitor of iNOS;

(c) the use of a PPAR delta agonist (e.g. a compound of formula (I)) in the manufacture of a medicament for the treatment of conditions for which an inhibitor of NO synthase and/or TNF is indicated, in particular, an inhibitor of iNOS;


(d) a method of treating conditions for which an inhibitor of NO synthase and/or TNF is indicated, in particular, an inhibitor of iNOS in a mammal, such as a human, which comprises the administration of a therapeutically effective amount of a PPAR delta agonist (e.g. a compound of formula (I)) to a said mammal.

The term 'treatment' as used herein includes prophylaxis as well as alleviation of established conditions for which an inhibitor of NO synthase and/or TNF is indicated. The PPAR delta agonists may be used as compounds or pharmaceutically acceptable derivatives thereof.

By pharmaceutically acceptable derivative is meant any pharmaceutically acceptable salt, solvate, ester or amide, or salt or solvate of such ester or amide, of the PPAR delta agonist, or any other compound which upon administration to the recipient is capable of providing (directly or indirectly) the PPAR delta agonist or an active metabolite or residue thereof.


Suitable PPAR delta agonists are described in for example WO 01/00603 which discloses compounds of formula (I), and pharmaceutically acceptable salts and, solvates thereof,

6




wherein:


- 5 X represents a COOH (or a hydrolysable ester thereof) or tetrazole group;
 X¹ represents NH, NCH₃, O, S, a bond (i.e. is absent), CH₂, or CH where the dashed line indicates that when X¹ is CH the depicted bond is a double bond;
 X² represents O or S;
 R¹ and R² independently represent H, CH₃, OCH₃, or halogen;
 10 n is 1 or 2;
 one of Y and Z is N and the other is S or O;
 y is 0, 1, 2, 3, 4 or 5;
 Each R³ independently represents CF₃ or halogen.
- 15 These compounds may be conveniently prepared by a general process wherein a moiety like A is coupled to an alcohol (B) using the Mitsunobu protocol (O. Mitsunobu, 1981 Synthesis, p 1) or by alkylation of A using a suitable non nucleophilic base such as K₂CO₃, Cs₂CO₃ or NaH, with an alkyl halide (C). Note that this synthesis is preferably carried out with the acid group protected by R.
- 20 Preferably, R is 1-6 alkyl which can be hydrolyzed off to give an acid of Formula (I), or if readily hydrolyzable, the resulting ester can be administered.



A



B




C

25


For example, when n is 1, Y is S, Z is N, and R³ is para-CF₃:

7




Intermediates of type A are commercially available. The synthesis of
 5 intermediates of type B is also illustrated below.

Furthermore, the tetrazole derivatives may be conveniently prepared by a
 general process wherein a moiety like D is coupled to an alcohol (B) using the
 Mitsunobu protocol (O. Mitsunobu, 1981 Synthesis, p 1), by alkylation of D using
 10 a suitable non nucleophilic base such as K₂CO₃, Cs₂CO₃ or NaH, with an alkyl
 halide (C) or by coupling of a moiety like E with an alkyl halide (C) using a
 suitable non nucleophilic base such as NaOH..



15 For example, when n is 1, Y is S, Z is N, and R³ is para-CF₃:

8




Preferred compounds of the invention are:

- 5 2-{2-methyl-4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxy}acetic acid
- 2-{2-methyl-4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-5-yl)methyl)sulfanyl]phenoxy}acetic acid
- 10 methyl 2-{4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxy}acetate
- 2-{4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxy}acetic acid
- (E)-3-[2-methyl-4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methoxy]phenyl]-2-propenoic acid
- 15 2-[3-chloro-4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenyl]acetic acid

A particularly preferred PPAR delta agonist is {2-methyl-4-[4-methyl-2-(4-trifluoromethyl phenyl) thiazol-5-ylmethylthio]phenoxy}-acetic acid and pharmaceutically acceptable salts, solvates, and hydrolyzable esters thereof (formula (II) below).

5



10 It will also be appreciated by those skilled in the art that the PPAR delta activator may also be utilized in the form of a pharmaceutically acceptable salt or solvate thereof. The physiologically acceptable salts include conventional salts formed from pharmaceutically acceptable inorganic or organic acids or bases as well as quaternary ammonium acid addition salts. More specific examples of suitable acid salts include hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, 15 perchloric, fumaric, acetic, propionic, succinic, glycolic, formic, lactic, maleic, tartaric, citric, palmoic, malonic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, fumaric, toluenesulfonic, methanesulfonic, naphthalene-2-sulfonic, benzenesulfonic hydroxynaphthoic, hydroiodic, malic, steroic, tannic and the like. Other acids such as oxalic, while not in themselves pharmaceutically acceptable, 20 may be useful in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable salts. More specific examples of suitable basic salts include sodium, lithium, potassium, magnesium, aluminium, calcium, zinc, N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, N-methylglucamine and procaine salts. References hereinafter to a PPAR delta activator include 25 both compounds and their pharmaceutically acceptable salts and solvates.

The PPAR delta activators and their pharmaceutically acceptable derivatives are 30 conveniently administered in the form of pharmaceutical compositions. Such compositions may conveniently be presented for use in conventional manner in

admixture with one or more physiologically acceptable carriers or excipients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

- 5 While it is possible that the PPAR delta activators may be therapeutically administered as the raw chemical, it is preferable to present the active ingredient as a pharmaceutical formulation.

10 The formulations include those suitable for oral, parenteral (including subcutaneous e.g. by injection or by depot tablet, intradermal, intrathecal, intramuscular e.g. by depot and intravenous), rectal and topical (including dermal, buccal and sublingual) or in a form suitable for administration by inhalation or insufflation administration although the most suitable route may depend upon for example the condition and disorder of the recipient. The
15 formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing into association the compounds ("active ingredient") with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into
20 association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

Formulations suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets (e.g. chewable tablets in particular for
25 paediatric administration) each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

30 A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a other conventional excipients such as binding agents, (for example, syrup, acacia, gelatin, sorbitol, tragacanth,
35

mucilage of starch, polyvinylpyrrolidone) or hydroxymethyl cellulose or hydroxymethyl cellulose fillers (for example, lactose, sugar, microcrystalline cellulose, maize-starch, calcium phosphate or sorbitol), lubricants (for example, magnesium stearate, stearic acid, talc, polyethylene glycol or silica),
5 disintegrants (for example, potato starch or sodium starch glycollate) or wetting agents, such as sodium lauryl sulfate. Moulded tablets may be made by moulding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active
10 ingredient therein. The tablets may be coated according to methods well-known in the art.

Alternatively, the compounds of the present invention may be incorporated into oral liquid preparations such as aqueous or oily suspensions, solutions,
15 emulsions, syrups or elixirs, for example. Moreover, formulations containing these compounds may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents such as sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin, hydroxyethylcellulose, carboxymethyl
20 cellulose, aluminum stearate gel or hydrogenated edible fats; emulsifying agents such as lecithin, sorbitan mono-oleate or acacia; non-aqueous vehicles (which may include edible oils) such as almond oil, fractionated coconut oil, oily esters, propylene glycol or ethyl alcohol; and preservatives such as methyl or propyl p-hydroxybenzoates or sorbic acid. Such preparations may also be formulated as
25 suppositories, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

Formulations for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats
30 and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents.

The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried
35 (lyophilised) condition requiring only the addition of a sterile liquid carrier, for

example, water-for-injection, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

- 5 Formulations for rectal administration may be presented as a suppository with the usual carriers such as cocoa butter, hard fat or polyethylene glycol.

- 10 Formulations for topical administration in the mouth, for example buccally or sublingually, include lozenges comprising the active ingredient in a flavoured basis such as sucrose and acacia or tragacanth, and pastilles comprising the active ingredient in a basis such as gelatin and glycerin or sucrose and acacia.

- 15 For topical administration to the epidermis, the compounds may be formulated as creams, gels, ointments or lotions or as a transdermal patch:

- 20 The compounds may also be formulated as depot preparations. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

- 25 For intranasal administration the compounds of the invention may be used, for example as a liquid spray, as a powder or in the form of drops.

- 30 For administration by inhalation the compounds according to the invention are conveniently delivered in the form of an aerosol spray presentation from pressurised packs or a nebuliser, with the use of a suitable propellant, e.g. 1,1,1,2-trifluoroethane (HFA 134A) and 1,1,1,2,3,3,3, - heptapropene (HFA 227), carbon dioxide or other suitable gas. In the case of a pressurised aerosol the dosage until may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

In addition to the ingredients particularly mentioned above, the formulations may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavouring agents.

5

It will be appreciated by those skilled in the art that reference herein to treatment extends to prophylaxis as well as the treatment of established diseases or symptoms. Moreover, it will be appreciated that the amount of the PPAR delta activator required for use in treatment will vary with the nature of the condition being treated and the age and the condition of the patient and will be ultimately at the discretion of the attendant physician or veterinarian. In general, however, doses employed for adult human treatment will typically be in the range of 0.02-5000 mg per day, preferably 1-1500 mg per day. The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example as two, three, four or more sub-doses per day. The formulations according to the invention may contain between 0.1-99% of the active ingredient, conveniently from 30-95% for tablets and capsules and 3-50% for liquid preparations.

10

15

20 The PPAR delta activators for use in the instant invention may be used in combination with one or more other therapeutic agents for example, pain relievers such as a glycine antagonist, a sodium channel inhibitor (e.g. lamotrigine), a substance P antagonist (e.g. an NK1 antagonist), acetaminophen or phenacetin; a matrix metalloproteinase inhibitor; a nitric oxide synthase (NOS) inhibitor (e.g. an iNOS or an nNOS inhibitor); an inhibitor of the release, or
25 action, of tumour necrosis factor α ; an antibody therapy (e.g. a monoclonal antibody therapy); a stimulant, including caffeine; an H₂-antagonist, such as ranitidine; a proton pump inhibitor, such as omeprazole; an antacid, such as aluminium or magnesium hydroxide; an antifatulent, such as simethicone; a
30 decongestant, such as phenylephrine, phenylpropanolamine, pseudoephedrine, oxymetazoline, epinephrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine; an antitussive, such as codeine, hydrocodone, carmiphen, carbetapentane, or dextromethorphan; a diuretic; or a sedating or non-sedating antihistamine; a COX-2 inhibitor such as VioxxTM or CelebrexTM
35 and other NSAIDs. The invention thus provides in a further aspect the use of a

combination comprising a compound of formula (I) with a further therapeutic agent in the treatment of iNOS and/or TNF mediated diseases.

5 When the PPAR delta activators are used in combination with other therapeutic agents, the compounds may be administered either sequentially or simultaneously by any convenient route.

10 The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical formulations comprising a combination as defined above optimally together with a pharmaceutically acceptable carrier or excipient comprise a further aspect of the invention. The individual components of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations.

15 When combined in the same formulation it will be appreciated that the two compounds must be stable and compatible with each other and the other components of the formulation and may be formulated for administration. When formulated separately they may be provided in any convenient formulation, 20 conveniently in such a manner as are known for such compounds in the art.

When a PPAR delta activator is used in combination with a second therapeutic agent active against the same disease, the dose of each compound may differ from that when the compound is used alone. Appropriate doses will be readily 25 appreciated by those skilled in the art.

The present inventors have found that, chemical compounds which are activators of PPAR delta are also useful in the treatment of diseases or conditions where an inhibitor of NO synthase and/or TNF is indicated, in particular iNOS. 30 The identification of novel PPAR delta activators may lead to more effective drugs for the treatment of these diseases and conditions. Accordingly, the invention further provides a method for screening for compounds which will be useful in treating diseases or conditions where an inhibitor of NO synthase and/or TNF is indicated comprising the step of 35 determining whether the compound interacts directly with PPAR delta, or the step

of determining whether the compound activates PPAR delta. Suitable screening tests include the binding and transfection assays discussed hereinabove and disclosed in WO 00 08002. A suitable control for PPAR delta is for example 2-{2-methyl-4-[(4-methyl-2-[3-fluoro-4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxy}acetic acid which conveniently may be radiolabelled.


The present invention also provides a method for treating diseases of conditions where an inhibitor of NO synthase and/or TNF is indicated, in particular iNOS in a subject comprising the administration of therapeutically effective amounts of compounds identified using the method of the invention. The invention also provides the use of a compound identified by a screening method of the invention in the manufacture of a medicament for the treatment of these diseases or conditions.

The following examples are set forth to illustrate the synthesis of some particular compounds of the present invention and to further exemplify particular applications of general processes described above. Accordingly, the following Example section is in no way intended to limit the scope of the invention contemplated herein.

Examples

Example 1


Preparation of 2-{2-methyl-4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxy}acetic acid



Intermediate A


Chlorosulfonic acid (15mL) was cooled to 0°C. then 10.0 g (0.05M) of ethyl (2-methylphenoxyacetate) was added over 10 min. The reaction mixture was stirred at 0-5°C for 30m, the bath was removed and stirring continued for

2h. The reaction mixture was poured into ice, forming a white solid which was washed with ice water and dried under high vacuum affording the title compound (12.846 g ,86%).



Intermediate B:


To a well stirred solution of LiAlH_4 (1.52 g, 40 mmol) in dry THF (50 mL) at 0°C , was slowly added a solution of ethyl 4-methyl-2-[4-(trifluoromethyl)phenyl]-thiazole-5-carboxylate (12.6 g, 40 mmol) in dry THF (50 mL). The mixture was stirred at room temperature for 2 hs. The reaction was quenched by slow addition at 0°C of water (2 mL), 5N NaOH (2 mL) and water (6 mL). The precipitate was filtered, washed with EtOAc, MeOH, CH_2Cl_2 and THF. After evaporation, a yellow solid was obtained, that was crystallized from MeOH-water to afford intermediate B depicted above (9.90 g, 36 mmol, 90%) as a yellow solid mp $120\text{--}122^\circ\text{C}$.



Intermediate C:

To a cold (0°C) stirred solution of intermediate B (8.2g, 30 mmol) and Et₃N (6.07 g, 8.36 mL, 60 mmol), in dry CH₂Cl₂ (120 mL) was slowly added MeSO₂Cl (5.49 g, 3.71mL, 48 mmol). After 2 hs at 0°C more Et₃N (6 mmol) and MeSO₂Cl (4.8 mmol) were added. After 2 more h a tlc (hexane:EtOAc, 1:1) showed complete reaction. The reaction mixture was diluted with CH₂Cl₂ (120 mL) and washed with NaHCO₃ (sat.) (2 x 240 mL) and water (2 x 240 mL), dried, filtered and evaporated to afford intermediate C (8.0 g, 27 mmol, 90%) as a yellow solid.

2-{2-methyl-4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxy}acetic acid:



Intermediate A (4.68g, 16mM) was refluxed with 9.6 g of tin powder in ethanol (20mL) and dioxane/HCl (20 mL). After 3 h the reaction mixture was poured into ice and CH₂Cl₂ (200mL) and filtered. The phases were separated and the aqueous layer was extracted 2X 50 mL CH₂Cl₂. The combined organic layers were dried (MgSO₄), filtered and evaporated to yield 3.5g (97%). This material readily forms disulfides and therefore was used immediately. It was dissolved in acetonitrile (50mL) with intermediate C (4.0 g, 14.0mM) and Cs₂CO₃ (10.1 g, 31.0 mM) and stirred for 1 h then diluted with ether (200mL) and water (200mL). The phases were separated and the organic phase was washed 2X NaOH 0.1N (50mL), dried (MgSO₄), filtered and evaporated to afford crude product (6.57 g,) which was slurried in hexane:ether (1:1) and filtered to yield pure intermediate D (5.0g, 74%). This material was hydrolyzed as described below to prepare the title compound. A solution of the corresponding ester (Intermediate D) (1 mmol) in THF (10 mL) (in some cases few drops of MeOH were added to help solubility), was treated with 1N LiOH in water (2 mL, 2 mmol), and stirred 16 h at room temperature (when reactions were slow, the temperature was elevated to 50°C). The solution was neutralized with 1N HCl (2 mL, 2 mmol) and the organic solvent evaporated to afford an aqueous solution with an insoluble product. If the insoluble was a solid, it was filtered and dried to afford the final product. If the insoluble was an oil, it was extracted with EtOAc (30 mL). The organic solution was separated, washed with water (2 x 30 mL), dried, filtered, and evaporated to afford the final product.


25

Example 2

2-{2-methyl-4-[[{4-methyl-2-[3-fluoro-4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxy}acetic acid

30


18



Intermediate 1


NaHCO₃ (2 mmol) was added to a solution of P₄S₁₀ (0.2 mmol) in toluene (100 mL) and the mixture heated to reflux for ca. 30min. The substituted benzamide (1 mmol) was added and the reaction stirred at 90°C for 1h. The reaction was then evaporated to dryness, treated with brine (100 mL) and extracted with CH₂Cl₂ (2 X 50 mL). The organic phase dried, filtered, and evaporated to afford the final product as a yellow solid (50%).

MS m/z 223



Intermediate 2


Intermediate 1 was reacted with ethyl-2-chloroacetoacetate (1:1 mmol) and the mixture heated to reflux overnight. The reaction was cooled to room temperature and the solvent evaporated. The solid was crystallised from Et₂O or hexane to afford the the title compound as an off-white solid (56%). MS m/z 333



Intermediate 3

Intermediate 2 was reacted with a solution of LiAlH₄ at 0°C. The reaction was stirred while it was allowed to warm to rt. After all the starting material had disappeared, the reaction was cautiously treated with water (5mL) followed by 1N NaOH (10mL). The mixture was filtered through celite. The filtrate was extracted with CH₂Cl₂ (3 x 50mL). The organic phase was dried, filtered and evaporated to afford the title compound as an off-white solid (63%).


MS m/z 291



Intermediate 4


To a solution of the Intermediate 3 (1 mmol) and Et₃N (2 mmol) in CH₂Cl₂ (100 mL) at 0°C was added dropwise methanesulfonyl chloride (1.6 mmol). After 2-4h the reaction was complete. CH₂Cl₂ (50 mL) is added and the organic phase washed with a saturated NaHCO₃ solution (2 x 50 mL), water (2 x 50 mL), dried, filtered and then evaporated to afford the title compound as a pale yellow solid (100%).

¹H-NMR (CDCl₃) δ2.40 (s, 3H), 4.70 (s, 2H), 7.55-7.75 (m, 3H).



Intermediate 5

Chlorosulfonic acid (15mL) was cooled to 0°C. then 10.0 g (0.05M) of ethyl (2-methylphenoxy)acetate was added over 10 m. The reaction mixture was stirred at 0-5°C for 30m, the bath was removed and stirring continued for 2 h. The reaction mixture was poured into ice, forming a white solid which was washed with ice water and dried under high vacuum affording the title compound (12.846 g, 86%).



Example 2a


ethyl 2-methyl-4-((4-methyl-2-[3-fluoro-4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxyacetate

Intermediate 5 (4.68g, 16mM) was refluxed with 9.6 g of tin powder in ethanol (20mL) and dioxane/HCl (20 mL). After 3 h the reaction mixture was poured into ice and CH₂Cl₂ (200mL) and filtered. The phases were separated and the aqueous layer was extracted 2X 50 mL CH₂Cl₂. The combined organic layers were dried (MgSO₄), filtered and evaporated to yield 3.5g (97%). This material readily forms disulfides and therefore was used immediately. It was dissolved in

acetonitrile (50mL) with intermediate 4 (4.0 g, 14.0mM) and Cs_2CO_3 (10.1 g, 31.0 mM) and stirred for 1 h then diluted with ether (200mL) and water (200mL). The phases were separated and the organic phase was washed 2X NaOH 0.1N (50mL), dried (MgSO_4), filtered and evaporated to afford crude product (6.57 g,)
5 which was intermediate 54. The crude material was chromatographed CH_2Cl_2 (100%) to afford the title compound (50%) as a clear oil that solidified on standing.

1H-NMR (CDCl_3) δ 1.20 (t, 3H), 2.15 (s, 3H), 2.20 (s, 3H), 4.05 (s, 2H), 4.15 (q, 2H), 4.55 (s, 2H), 6.55 (d, 1H), 7.05 (dd, 1H), 7.15 (d, 1H), 7.55 (t, 1H), 7.65 (m, 2H).
10

MS m/z 500 (M+1)



Example 2b

2-{2-methyl-4-[(4-methyl-2-[3-fluoro-4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl]sulfanyl]phenoxy}acetic acid
15

A solution of Example 2a (1 mmol) in THF (10 mL) was treated with 1N LiOH in water (2 mL, 2 mmol), and stirred 16 h at room temperature (when reactions were slow, the temperature was elevated to 50°C). The solution was neutralized with 1N HCl (2 mL, 2 mmol) and the organic solvent evaporated to afford an aqueous solution with an insoluble product. If the insoluble was a solid, it was
20 filtered and dried to afford the final product. If the insoluble was an oil, it was extracted with EtOAc (30 mL). The organic solution was separated, washed with water (2 x 30 mL), dried, filtered, and evaporated to afford the final product.

25

The crude material was precipitated from ether to afford the title compound (81%) as a yellow foam: mp <50°C.

Anal. Calcd. for $\text{C}_{21}\text{H}_{17}\text{F}_4\text{NO}_3\text{S}_2$: C, 53.50; H, 3.63; N, 2.97; S, 13.60. Found: C53.86; H, 3.63; N, 2.87; S, 13.82.

30

MS m/z 472 (M+1)

INOS biology

The ability of PPAR delta activators to inhibit the activity and expression of inducible nitric oxide synthase (INOS) was examined using potent, selective PPAR delta agonists, such as (2-{2-methyl-4-[(4-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxy}acetic acid (hereinafter referred to as Compound A, synthesis as described in Example 1) and (2-{2-methyl-4-[(4-methyl-2-[3-fluoro-4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl)sulfanyl]phenoxy}acetic acid (hereinafter referred to as Compound B, synthesis as described in Example 2b).

In this assay system, lipopolysaccharide (LPS) was used to induce the expression of INOS in the mouse macrophage cell line, J774. Pre-treatment of J774 cells, with pharmacologically relevant concentrations of PPAR delta agonists, resulted in inhibition of the INOS activity. This reduction in INOS activity correlated with a reduction in LPS-induced INOS mRNA expression.

Measurement of iNOS Activity

Specifically, LPS-induced INOS activity was measured using the following assay conditions. J774 cells were seeded at a density of 35000-50000 thousand cells per well, in a black, clear-bottomed, 96 well plate, 24 hours prior to use. The cell culture and the drug dilutions were carried out in complete media, which consisted of DMEM (Dulbecco's modification of Eagle's medium) containing foetal calf serum (10%), glutamine (2mM), penicillin (100u/ml) and streptomycin (100µg/ml). The J774 cells were pre-treated with the PPAR delta activators A and B, or vehicle, for 6 hours prior to, and for 24 hours subsequent to the addition of LPS (typically, a concentration of 1ug/ml of LPS was used). Twenty-four hours after the addition of LPS, INOS activity was measured by the following method. The cell culture media/drug dilutions were removed and the cells washed with D-PBS (Dulbecco's modification of phosphate-buffered saline). The D-PBS was then removed, and replaced with D-PBS containing DAF-2 (4,5-diaminofluorescein ; 5uM) and L-arginine (500uM). After incubation at 37°C for 3 hours, fluorescence from each well was measured at an excitation wavelength of 485nm, and an emission wavelength of 530nm. The ability of

LPS to induce INOS activity, in the presence and absence of PPAR delta activators, was then calculated. Examples of the data obtained are shown in Figure 1, with the control elevation of INOS activity, induced by LPS, is a response of 100%.

5

Thus, pre-treatment with Compound A or Compound B, produced a marked inhibition of LPS-induced INOS activity in J774 cells, at sub-nanomolar – nanomolar concentrations.

10 Measurement of Inhibition of INOS mRNA

The mechanism, associated with the inhibition of INOS activity by PPAR delta activators, was investigated in greater detail.

Specifically, the ability of PPAR delta agonists to inhibit LPS-induced expression of INOS mRNA, in J774 cells was examined. J774 cells were plated in 6 well plates (10^6 cells/well), 24 hours prior to use. The cells were pre-treated with PPAR delta activator control media for 6 hours, prior to addition of LPS, which was co-incubated with the PPAR delta activator / control, for a further 24 hours. At the end of this incubation period, the culture medium was removed by aspirating and the cells washed with D-PBS. Following removal of the D-PBS, total cellular RNA was isolated from each sample using a commercially available RNA isolation kit. First strand cDNA synthesis was carried out as per instructions supplied with the AMV reverse transcription (RT) system. An aliquot (1000ng) of the RNA was added to a mix which contained (final concentrations) 15 MgCl₂ (5mM), Tris-HCl (10mM ; pH 8.8), KCl (50mM), Triton X-100 (0.1%), 25 dNTP (1mM), rNasin (1U/ μ l), AMV reverse transcriptase (0.75U/ μ l), oligo(dT)₁₅ (25ng/ μ l). This was incubated in a thermal cycler at 42°C for 30 minutes, followed by 95°C for 15 minutes, and, finally, 4°C, until being transferred to a freezer (-20°C) for storage.

30

For use in PCR, the following primer sets were used

Mouse INOSsense	CCCTTCCGAAGTTTCTGGCAGCAGC (SEQ ID NO: 1)
Mouse INOS anti-sense	GGCTGTCAGAGCCTCGTGGCTTTGG (SEQ ID NO: 2)
Mouse GAPDH sense	TGAAGGTCGGTGTGAACGGATTTGGC (SEQ ID NO: 3)
35 Mouse GAPDH anti-sense	CATGTAGGCCATGAGGTCCACCAC (SEQ ID NO: 4)

PCR was undertaken in a 50µl reaction volume containing 5µl of the RT reaction, sense and anti-sense primers for INOS/GAPDH (0.4pmol/µl), dNTPs (160µM), KCl (50mM), Tris-HCl (10mM ; pH 9.0), Triton X-100 (0.1%), MgCl₂ (2mM), and Taq DNA polymerase (0.04U/µl) (final concentrations). The PCR was carried out in a thermal cycler using the following conditions : 95°C for 60s, followed by 28 cycles of 94°C for 30s, 55°C for 60s, 72°C for 90s. Following a final extension step of 72°C for 5 minutes, the samples were maintained at 4°C, until analysed on an agarose gel. Analysis of sybr-green stained gels, by densitometry, was carried out using a Storm fluorimager system (Molecular Devices)

Thus, pre-treatment of J774 cells, with pharmacologically relevant concentrations of PPAR delta activators, resulted in inhibition of LPS(1µg/ml for data below)-induced INOS mRNA expression. This data is shown in Figure 2.

Measurement of Inhibition of TNF


The ability of PPAR delta activators to inhibit the expression/secretion of tumour necrosis factor-α (TNF) was also examined. Thus, the ability of PPAR delta activators to inhibit LPS- induced TNF expression correlated well with the inhibitory effects of PPAR delta activators on LPS-induced INOS activation.

Specifically, LPS-induced INOS activity and expression of TNF was measured using the following assay conditions. J774 cells were seeded at a density of 35000-50000 thousand cells per well, in black, clear-bottomed, 96 well plates. The cell culture and the drug dilutions were carried out in complete media, which consisted of DMEM (Dulbecco's modification of Eagle's medium) containing foetal calf serum (10%), glutamine (2mM), penicillin (100u/ml) and streptomycin (100µg/ml). The J774 cells were pre-treated with PPAR delta activators, or vehicle, for 6 hours prior to, and for 24 hours subsequent to the addition of LPS (typically, a concentration of 1µg/ml of LPS was used). Twenty-four hours after the addition of LPS, INOS activity was measured by the following method. The cell culture media/drug dilutions were removed for measurement of TNF concentrations, quantified using a commercially available ELISA system. The cells washed with D-PBS. The D-PBS was then removed, and replaced with D-

PBS containing DAF-2 (4,5-diaminofluorescein ; 5 μ M) and L-arginine (500 μ M). INOS activity was then measured as described previously. Examples of the data obtained are illustrated in Figure 3.

Claims

1. A method of treating diseases or conditions for which an inhibitor of NO synthase and/or TNF is indicated in a mammal, such as a human, which comprises the administration of a PPAR delta activator.
2. Use of a PPAR delta activator in the manufacture of a medicament for the treatment of diseases or conditions for which an inhibitor of NO synthase and/or TNF is indicated.
3. A method according to use according to claim 1 or use according to claim 2 wherein the PPAR delta activator is a PPAR delta agonist.
4. A method of use according to claim 3 wherein the PPAR delta agonist is a selective agonist.
5. A method of use according to claim 4 wherein the selective PPAR delta agonist is a hPPAR delta selective agonist.
6. Use according to claim 2 or a method according to claim 1 where the PPAR delta activator has a structure according to formula (I).



7. Use or a method according to claim 5 wherein the PPAR delta activator is {2-methyl-4-[4-methyl-2-(4-trifluoromethyl phenyl) thiazol-5-ylmethylthio]phenoxy}-acetic acid.

8. A pharmaceutical formulation for use in the treatment of diseases or conditions for which an inhibitor of NO synthase and/or TNF is indicated comprising a PPAR delta activator together with at least one pharmaceutical carrier wherein the PPAR delta agonist is present in an amount effective for use in the treatment of diseases or conditions for which an inhibitor of NO synthase and/or TNF is indicated.
5
9. A method for identifying compounds which will be useful in treating diseases or conditions for which an inhibitor of NO synthase and/or TNF is indicated comprising the step of determining whether the compound interacts directly with PPAR delta, or the step of determining whether the compound activates PPAR delta.
10
10. The present invention also provides a method for treating diseases or conditions for which an inhibitor of NO synthase and/or TNF is indicated comprising administration of compounds identified using the screening method of claim 9
15

Figure 1





Figure 2






Figure 3



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 April 2002 (11.04.2002)

PCT

(10) International Publication Number
WO 02/028433 A3

(51) International Patent Classification⁷: A61K 45/08,
31/426, 31/421, A61P 1/00, 19/02

(21) International Application Number: PCT/GB01/04370

(22) International Filing Date: 2 October 2001 (02.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0024362.6 5 October 2000 (05.10.2000) GB

(71) Applicant (for all designated States except US): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BUCHAN, Kevin, William** [GB/GB]; c/o GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB).

(74) Agent: **GIDDINGS, Peter, John**; GlaxoSmithKline, Corporate Intellectual Property (CN9.25.1), 980 Great West Road, Brentford, Middlesex TW8 9GS (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
15 August 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/028433 A3

(54) Title: USE OF A PPAR DELTA ACTIVATOR IN THE TREATMENT OF DISEASES RELATED TO NO INHIBITION OR TNF INHIBITION

(57) Abstract: Methods of prevention or treatment of diseases or conditions where inhibition of NO synthase and/or TNF is desirable, the use of PPAR delta activators in such methods and methods for the identification of compounds useful in such treatment.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 01/04370

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K45/08 A61K31/426 A61K31/421 A61P1/00 A61P19/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, CHEM ABS Data, PASCAL, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
L	WO 01 00603 A (SIERRA MICHAEL LAWRENCE ;GELLIBERT FRANCOISE JEANNE (FR); GLAXO GR) 4 January 2001 (2001-01-04) cited in the application claims 24-26	2,8
X	WO 97 28149 A (AUWERX JOHAN ;BERGER JOEL P (US); MERCK & CO INC (US); MOLLER DAVI) 7 August 1997 (1997-08-07) claim 54	2,8
X	EP 1 023 907 A (YAMANOUCHI PHARMA CO LTD) 2 August 2000 (2000-08-02) * p,3, [0012] *	2,8
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

21 March 2002

Date of mailing of the international search report

10 7. 06. 02

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Uiber, P

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 01/04370

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE MEDLINE [Online] NLM9793108, 1998 CITROME ET AL: "Utilization of valproate: extent of inpatient use in the New York State Office of Mental Health." XP002193787 abstract & THE PSYCHIATRIC QUARTERLY., vol. 69, no. 4, 1998, pages 283-300, ---	2,8
P,X	EP 1 170 008 A (CHEMOTHERAPEUTISCHES FORSCHUNG) 9 January 2002 (2002-01-09) * p.3, 1.15; claims 1-15 * -----	2,8

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 01/04370

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1, 3-5, 10
because they relate to subject matter not required to be searched by this Authority, namely:
Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy (claims 1, 3-5, 10)
2. ☒ Claims Nos.: 2, 8
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
2 (part), 6, 7, 8 (part)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 2,8

Present claims 2 and 8 relate to a pharmaceutical product or a method respectively containing or testing a compound defined by reference to a desirable characteristic or property, namely PPAR delta activators. In addition, the diseases or conditions are defined by reference to a desirable characteristic or property, namely, " for which an inhibitor of NO synthetase and/or TNF is indicated".

The claims cover all a pharmaceutical product or a method respectively containing or testing SAID compound having this characteristic or property or for treating said diseases or conditions, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such a pharmaceutical product, a method respectively containing or testing a compound or such diseases or conditions.

In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product and the diseases or conditions by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds of formula (I) in claim 6 for treating the diseases EXPLICITLY mentioned in pages 2-4.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 2(part.),6,7,8(part.)

Use of a PPAR delta activator of formula I in the manufacture of a medicament for treating diseases or conditions for which an inhibitor of NO synthetase and/or TNF is indicated and pharmaceutical compositions containing said PPAR delta activators

2. Claims: 9,10

A method for identifying a PPAR delta activator not falling within formual I and its use in the manufacture of a medicament for treating diseases or conditions for which an inhibitor of NO synthetase and/or TNF is indicated

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/04370

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0100603	A	04-01-2001	AU 5817100 A	31-01-2001
			BR 0011891 A	05-03-2002
			CZ 20014664 A3	13-03-2002
			WO 0100603 A1	04-01-2001
			EP 1189895 A1	27-03-2002
			NO 20016078 A	13-12-2001

WO 9728149	A	07-08-1997	AU 1856997 A	22-08-1997
			AU 721452 B2	06-07-2000
			AU 2115997 A	22-08-1997
			CA 2245529 A1	07-08-1997
			EP 0888278 A1	07-01-1999
			JP 2002503202 T	29-01-2002
			WO 9728115 A1	07-08-1997
			WO 9728149 A1	07-08-1997
			AU 712607 B2	11-11-1999
			AU 1858197 A	22-08-1997
			CA 2244831 A1	07-08-1997
			EP 1011651 A1	28-06-2000
			JP 2000504021 T	04-04-2000
			WO 9727847 A1	07-08-1997
			AU 719146 B2	04-05-2000
			AU 2250797 A	22-08-1997
			CA 2245524 A1	07-08-1997
			EP 0904079 A1	31-03-1999
			WO 9727857 A1	07-08-1997
			AU 708055 B2	29-07-1999
			AU 1856397 A	22-08-1997
			EP 0882029 A1	09-12-1998
			JP 2002503203 T	29-01-2002
			WO 9728137 A1	07-08-1997
			US 5859051 A	12-01-1999
			US 6090836 A	18-07-2000
			ZA 9700824 A	30-10-1998
			US 6020382 A	01-02-2000
			US 5847008 A	08-12-1998
			AU 719663 B2	11-05-2000
			AU 5615298 A	17-07-1998
			EP 0948327 A1	13-10-1999
			JP 2001511767 T	14-08-2001
			WO 9827974 A1	02-07-1998
			US 6160000 A	12-12-2000
			US 6090839 A	18-07-2000

EP 1023907	A	02-08-2000	AU 8355998 A	16-02-1999
			EP 1023907 A1	02-08-2000
			US 6300364 B1	09-10-2001
			WO 9904815 A1	04-02-1999

EP 1170008	A	09-01-2002	EP 1170008 A1	09-01-2002
			WO 0207722 A2	31-01-2002

THIS PAGE BLANK (USPTO)